

## Trans-Labrador Sea Modern Reflection Data Show Unorthodox Rift

Michael E. Enachescu\*  
Memorial University, St. John's, NL, Canada  
michaele@mun.ca

and

Paul H. Einarsson  
Geophysical Service Inc., Calgary, AB, Canada

During 2006 Geophysical Service Incorporated (GSI) acquired two regional multi-channel seismic (MCS) reflection lines across the Labrador Sea at the approximate latitude of the northern Saglek Basin, on the Labrador shelf (Canada) and Lady Franklin Basin on the Greenland shelf (Denmark). More precisely, the two NE-SW trending lines run between 59° and 63° North Latitude and 51° to 63° West Longitude and are each approximately 750 km long. The data was acquired using a 7200 m streamer, a 37.5 m source point interval, and a 12.5 m group interval resulting in a nominal fold of ninety-six. The lines were recorded to 12 seconds and processed to prestack time migration using a Kirchhoff curved ray algorithm.

These data are crucial for deciphering the nature of the basement beneath the sedimentary cover of the Labrador Sea shelf, slope and rise. They also help to clarify the evolution of the Labrador Sea and its sedimentary basins through the following stages: 1) Mesozoic intra-cratonic extension (intercontinental rifting); 2) mantle exhumation and transitional crust formation; 3) Cenozoic oceanic crust creation (oceanic rifting) and 4) cessation of oceanic rifting (drifting) with associated prominent regional thermal subsidence. The lines are complementary to earlier MCS research data in the area and offer critical information in determining:

- (a) if any true oceanic crust was emplaced between the northern Labrador and Greenland margins;
- (b) the location, timing, extent and modality of such emplacement; and
- (c) if a continuous mid-ocean ridge was active in the area.

These lines will also provide fundamental geoscientific data in support of Canada's ongoing United Nations Convention on the Law of the Sea (UNCLOS) negotiations regarding the delineation of maritime jurisdictional boundaries in the Labrador Sea. Of particular interest is that the lines show the presence of thick Mesozoic and probably older sedimentary basins with potential hydrocarbon resources, in currently drillable water depths (3km) lying far seaward of the 200 NM limit, and well beyond any boundary that might be defined by currently proposed bathymetric criteria. In this regard the GSI lines can be used to support an extension of Canada's

jurisdiction boundary to include new areas of significant environmental and economic interest. Similar observations relating to the extension of sedimentary basins into the Labrador Sea deepwater have also been made using data sets that were recorded for petroleum exploration on the Greenland margin.

#### **Acknowledgements**

Geophysical Service Inc., Pan-Atlantic Petroleum Systems Consortium (PPSC), NSERC, PRAC

#### **Further Reading**

Balkwill, H.R., McMillan, N. J., MacLean, B., Williams, G.L. & Srivastava, S.P., 1990. Geology of the Labrador Shelf, Baffin Bay and Davis Strait. In: Keen, M. J. & Williams, G. L. (eds) Geology of the continental margins of eastern Canada. Geology of Canada, 2, 293–348.

Chalmers, J.A. and Laursen, K.H., 1995. Labrador Sea: the extent of continental crust and the timing of the start of sea-floor spreading. *Marine and Petroleum Geology*, 12, 205–217.

Chalmers, J.A. and Pulvertaft, T.C.R., 2001. Development of the continental margins of the Labrador Sea – a review. In: Wilson, R.C.L., Whitmarsh, R.B., Taylor, B. & Froitzheim, N. (eds) Non-Volcanic Rifting of Continental Margins: A comparison of Evidence from Land and Sea. Geological Society, London, Special Publications 187, 77–105.

Commission on the Limits of the Continental Shelf (CLCS), 1999. Scientific and technical guidelines of the Commission on the Limits of the Continental Shelf. CLCS/11, 91 pp. + annexes.

Enachescu, M., 2006. Hopedale Basin-1: Favorable geology, advanced technology may unlock Labrador's substantial resources, *Oil and Gas Journal*, 19 June, v. 104.23, p. 29-34.

Enachescu, M., 2006. Hopedale Basin-2: Atlantic of Labrador poised for modern exploration round, *Oil and Gas Journal*, 26 June, v. 104.24, p. 36-42.

Enachescu, M., J. Hogg and S. Kearsey, 2006. The Hopedale Basin, Offshore Labrador, Canada: Stranded Gas and Remaining Petroleum Potential, AAPG Convention, Houston, Tx.

Srivastava, S. P., Falconer, R. K. H. and MacLean, B., 1981. Labrador Sea, Davis Strait, Baffin Bay: Geology and Geophysics – A Review. In *Geology of the north Atlantic Borderlands*, eds. J. Wm. Kerr and A.J. Fergusson. Can. Petrol. Geol. Memoir 7, 333 – 398.

Srivastava, S. P., Verhoef, J. and Macnab, R., 1988. Results from a detailed aeromagnetic survey across the northeast Newfoundland margin, Part I: Spreading anomalies and relationship between magnetic anomalies and the ocean-continent boundary. *Marine and Petroleum Geology* 5, 306-323.

Srivastava, S. P., and C. E. Keen, 1995. A deep seismic reflection profile across the extinct mid-Labrador Sea spreading center, *Tectonics*, 14(2), 372–389.