

# Functional Gridding of Geophysical data for high resolution mapping of local anomalies

Fatih Doğan, PhD. Physics  
TerraNotes Ltd. Edmonton AB  
[fatih.terrannotes@gmail.com](mailto:fatih.terrannotes@gmail.com)

Volkan Tuncer, M.Sc., P.Geoph.  
TerraNotes Ltd, Edmonton AB  
[Volkan.terrannotes@gmail.com](mailto:Volkan.terrannotes@gmail.com)  
Lei Sha, M.Sc., P. Geoph.  
TerraNotes Ltd. Edmonton AB  
[lei.terrannotes@gmail.com](mailto:lei.terrannotes@gmail.com)

Julien Lambert, B.A.; GDM  
Terrannotes Ltd. Edmonton AB  
[terrannotes@gmail.com](mailto:terrannotes@gmail.com)

## Summary

I will present a method that we researched and developed to regularize geophysical data to a lattice using an accurate and fast functional gridding mechanism. I will present examples that demonstrate how this method can accurately interpolate irregular data. This method is especially useful to identify very localized structures, such as Kimberlite pipes.

## Introduction

In order to achieve high-resolution imaging, geophysical data must be processed from highly accurate, spatially regularized lattice.

## Theory and Method

Higher accuracy of gridded data improves the signal to noise ratio. We developed a gridding technique that builds a regular grid from randomly located points on 2D or 3D data sets. We used an iterative local update scheme to make the approximation of the data more accurate. As well, we utilized a mixture of global and local updates, decided by frequency of improvements on the approximation by each iteration step. The geological features present in the study area are assumed to be local and continuous through functional interpolation. We compared our results to several gridding techniques used in commercially available software, including Krigging, triangulation, radial basis, inverse distance to a power and minimum curvature.

## **Conclusions**

Our iterative local update technique has proven to be more accurate for the same or improved efficiency level. This technique can be applied to any spatial axis and is efficient enough to apply to large 3D surveys.

## **Acknowledgements**

TerraNotes and I would like to acknowledge the research and development contribution for this study from the following agencies:

1. National Research Council Canada (IRAP)
2. Alberta Ingenuity Fund (now Alberta Innovates Technology Futures)

And finally, we would like to thank to GeoScience 2010 for accepting our abstract and giving us the opportunity to share our study's findings with you.

## **References**

- Stein, M.L. (1999), Statistical Interpolation of Spatial Data: Some Theory for Kriging, Springer, New York.
- Smith, Richard S. and O'Connell, Michael D. Geophysics 2005, vol. 70, no 5, pp. V121-V127